

**36-3333: Anti-Tyrosinase-Related Protein-1 (TRP-1) (Melanoma Marker) Monoclonal Antibody(Clone: SPM456)**

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	SPM456
<b>Application :</b>	FACS,IF
<b>Reactivity :</b>	Human, Mouse
<b>Gene :</b>	TYRP1
<b>Gene ID :</b>	7306
<b>Uniprot ID :</b>	P17643
<b>Alternative Name :</b>	5, 6 dihydroxyindole 2 carboxylic acid oxidase, 6-dihydroxyindole-2-carboxylic acid oxidase, Associated with iris pigmentation, CAS2, Catalase B (CATB), DHICA oxidase, Glycoprotein75 (GP75), Melanoma antigen gp75, Tyrosinase-related protein 1 (TYRP1), TYRRP
<b>Isotype :</b>	Mouse IgG2a, kappa
<b>Immunogen Information :</b>	SK-MEL-23 cells

**Description**

It reacts with a 75kDa melanocyte-specific gene product, identified as Tyrosinase-related protein-1 (TRP-1). It is involved in melanin synthesis. TRP1 is present on the melanosomal membranes of melanoma, normal melanocytes and nevi. Recent evidence suggests that TRP-1 is involved in maintaining stability of tyrosinase protein and modulating its catalytic activity. TRP-1 is also involved in maintenance of melanosome ultrastructure and affects melanocyte proliferation and cell death.

**Product Info**

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

**Application Note**

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml);