

## 36-3410: Anti-TCL1 (T-Cell Marker) Monoclonal Antibody(Clone: TCL1/2747R)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	TCL1/2747R
<b>Application :</b>	IHC
<b>Reactivity :</b>	Human
<b>Gene :</b>	TCL1A
<b>Gene ID :</b>	10232
<b>Uniprot ID :</b>	P56279
<b>Alternative Name :</b>	TCL1A; Lymphoma/leukemia, T-cell; Oncogene TCL1; P14 TCL1; T cell leukemia 1; T cell lymphoma 1; T cell lymphoma 1A;T-cell leukemia/lymphoma 1A; TCL1; TCL1A
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Recombinant human TCL1 protein fragment (around aa 2-109) (exact sequence is proprietary)

### Description

T-cell leukemia/Lymphoma Protein 1A (TCL-1A), also known as p14TCL1 is a product of the TCL1 gene that is involved in T-cell prolymphocytic leukemia (TPLL). T-PLL is a rare form of mature T-cell leukemia, which is consistently associated with chromosomal rearrangements characterized by the juxtaposition of the TCRA locus on chromosome 14q11 and the TCL1A gene on 14q32.13. TCL1 is overexpressed in Burkitt lymphoma, the majority of AIDS-related non-Hodgkin lymphoma-designated immunoblastic plasmacytoid lymphoma, lymphoblastic lymphoma, chronic lymphocytic leukemia, mantle cell lymphoma, follicular lymphoma, diffuse large B-cell lymphoma, and primary cutaneous B-cell lymphoma.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Tonsil stained with TCL1 Recombinant Rabbit Monoclonal Antibody (TCL1/2747R).




Fig. 2: Formalin-fixed, paraffin-embedded human Tonsil stained with TCL1 Recombinant Rabbit Monoclonal Antibody (TCL1/2747R).




Fig. 3: SDS-PAGE Analysis Purified TCL1 Recombinant Rabbit Monoclonal Antibody (TCL1/2747R). Confirmation of Purity and Integrity of Antibody