

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-3488: Anti-CD4 (T-Helper/Inducer Cell Marker) Monoclonal Antibody(Clone: RIV7)

Clonality: Monoclonal

Clone Name: RIV7

Application : Functional Assay, FACS, IF

Reactivity: Human
Gene: CD4
Gene ID: 920
Uniprot ID: P01730

Alternative Name: L3T4; Leu3; Ly-4; Lymphocyte antigen CD4; p55; T cell antigen T4/LEU3; T cell differentiation

antigen L3T4; T-cell surface antigen T4/Leu-3; T-cell surface glycoprotein CD4

Isotype: Mouse IgG2a, kappa

Immunogen Information: Human peripheral lymphocytes

Description

Recognizes a protein of 55kDa, identified as CD4. It is a membrane glycoprotein of T lymphocytes that interacts with major histocompatibility complex class II antigens and is also a receptor for the human immunodeficiency virus. This protein is expressed not only in T lymphocytes, but also in B cells, macrophages, and granulocytes. It is also expressed in specific regions of the brain. The protein functions to initiate or aµgment the early phase of T-cell activation, and may function as an important mediator of indirect neuronal damage in infectious and immune-mediated diseases of the central nervous system. Multiple alternatively spliced transcript variants encoding different isoforms have been identified.

Product Info

Amount : 20 μg / 100 μg

Content: 200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS

with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage condition : Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous.

Application Note

Functional Studies (Order Ab without azide);, Flow Cytometry (1-2ug/million cells in 0.1ml); , Immunofluorescence (1-2ug/ml) ,




Fig. 1: SDS-PAGE Analysis Purified CD4 Mouse Monoclonal Antibody (RIV7). Confirmation of Purity and Integrity of Antibody.