

36-3493: Anti-CD5 (Mantle Cell Lymphoma Marker) Monoclonal Antibody(Clone: B-B8)

Clonality :	Monoclonal
Clone Name :	B-B8
Application :	FACS,IF
Reactivity :	Human
Gene :	CD5
Gene ID :	921
Uniprot ID :	P06127
Alternative Name :	CD5 antigen (p56 62), LEU1, Ly12, LyA, Lymphocyte antigen T1/Leu-1, Lymphocyte glycoprotein T1/Leu1, T-cell surface glycoprotein CD5
Isotype :	Mouse IgG1, kappa
Immunogen Information	: T cells activated with PHA

Description

Recognizes a 67kDa transmembrane protein which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas. Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplastic B-cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL), mantle cell lymphoma, and a subset (~10%) of diffuse large B-cell lymphoma. CD5 aberrant expression is useful in making a diagnosis of mature T-cell neoplasms. Note that this MAb is not suitable for frozen tissues.

Product Info

Amount :	20 μg / 100 μg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml);

		NR	R	
250→				
150→	-	-		
100→			à.	
75→		, ý		
50→			-	2 ug loading NR=Non-
37→				reduced R=reduced
25→	_		-	
20→				
15→				
10→				

Fig. 1: SDS-PAGE Analysis Purified CD5 Mouse Monoclonal Antibody (Clone B-B8). Confirmation of Integrity and Purity of Antibody.