∗ abeomics

36-3562: Anti-CD28 Monoclonal Antibody(Clone: C28/77)

| Clonality : | Monoclonal |
|-----------------------|---|
| Clone Name : | C28/77 |
| Application : | ELISA |
| Reactivity : | Human |
| Gene : | CD28 |
| Gene ID : | 940 |
| Uniprot ID : | P10747 |
| Alternative Name : | T-cell-specific surface glycoprotein CD28; Tp44 |
| Isotype : | Mouse IgG1, kappa |
| Immunogen Information | Recombinant human CD28 protein |

Description

Recognizes a glycoprotein of 44-88kDa, which is identified as CD28. It is the critical T-cell co-stimulatory receptor which provides to the cell the important second activation signal by binding CD80 and CD86 that are expressed by antigen presenting cells. Besides its co-stimulation role, CD28 functions in preventing T-cells from anergic hyporesponsive state or from undergoing premature apoptotic cell death. CD28 is also expressed on human fetal NK cells and some NK cell lines, whereas on murine NK cells the CD28 expression is much broader.

Product Info

| Amount : | 20 µg / 100 µg |
|---------------------|--|
| Content : | 200 μ g/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Storage condition : | Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. |

Application Note

ELISA (For coating, order Ab without BSA);




Fig. 1: SDS-PAGE Analysis Purified CD28 Mouse Monoclonal Antibody (C28/77). Confirmation of Integrity and Purity of Antibody.