

36-3585: Anti-CD33 / SIGLEC3 (Myeloid Cell Surface Antigen) Monoclonal Antibody(Clone: WM53)

| | |
|--------------------------------|---------------------------------------------------------------------------------------------------|
| Clonality : | Monoclonal |
| Clone Name : | WM53 |
| Application : | FACS,IF,Functional Assay |
| Reactivity : | Human |
| Gene : | CD33 |
| Gene ID : | 945 |
| Uniprot ID : | P20138 |
| Alternative Name : | gp67; My9; Myeloid cell surface antigen CD33; p67; Sialic acid-binding Ig-like lectin 3 (SIGLEC3) |
| Isotype : | Mouse IgG1, kappa |
| Immunogen Information : | Human AML cells |

Description

Recognizes a 67kDa glycoprotein, which is identified as CD33 (HLDA IV; WS Code M-505). CD33 is a transmembrane protein of the sialic acid-binding immunoglobulin-like lectin (Siglec) family. It belongs to the immuno-receptor tyrosine-based inhibitory motif (ITIM)-containing molecules able of recruiting protein tyrosine phosphatases SHP-1 and SHP-2 to signal assemblies; these ITIMs are also used for ubiquitin-mediated removal of the receptor from the cell surface. CD33 is expressed on cells of myelomonocytic lineage, binds sialic acid residues in N- and O-glycans on cell surfaces, and is a therapeutic target for acute myeloid leukemia. CD33 is expressed on myeloid progenitors, monocytes, granulocytes, dendritic cells and mast cells. It is absent on platelets, lymphocytes, erythrocytes and hematopoietic stem cells.

Product Info

| | |
|----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Amount : | 20 µg / 100 µg |
| Content : | 200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Storage condition : | Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. |

Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Functional Studies (Order Ab without BSA & Azide);