

## 36-3586: Anti-CD33 / SIGLEC3 (Myeloid Cell Surface Antigen) Monoclonal Antibody(Clone: WM53)-PE

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	WM53
<b>Application :</b>	FACS,IF
<b>Reactivity :</b>	Human
<b>Conjugate :</b>	PE
<b>Gene :</b>	CD33
<b>Gene ID :</b>	945
<b>Uniprot ID :</b>	P20138
<b>Alternative Name :</b>	gp67; My9; Myeloid cell surface antigen CD33; p67; Sialic acid-binding Ig-like lectin 3 (SIGLEC3)
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Human AML cells

### Description

Recognizes a 67kDa glycoprotein, which is identified as CD33 (HLDA IV; WS Code M-505). CD33 is a transmembrane protein of the sialic acid-binding immunoglobulin-like lectin (Siglec) family. It belongs to the immunoreceptor tyrosine-based inhibitory motif (ITIM)-containing molecules able of recruiting protein tyrosine phosphatases SHP-1 and SHP-2 to signal assemblies; these ITIMs are also used for ubiquitin-mediated removal of the receptor from the cell surface. CD33 is expressed on cells of myelomonocytic lineage, binds sialic acid residues in N- and O-glycans on cell surfaces, and is a therapeutic target for acute myeloid leukemia. CD33 is expressed on myeloid progenitors, monocytes, granulocytes, dendritic cells and mast cells. It is absent on platelets, lymphocytes, erythrocytes and hematopoietic stem cells.

### Product Info

<b>Amount :</b>	0.5 ml at 100µg/ml
<b>Content :</b>	Antibody Purified from Bioreactor Concentrate by Protein A/G and conjugated to various reporter molecules. Prepared in 10mM PBS with 0.05% BSA and 0.05% azide. Contact us if you require this Ab in a different format.
<b>Storage condition :</b>	Antibody with azide - store at 4 to 8°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Flow Cytometry: 5ul per test per one million cells (or 5ul per 100ul of whole blood);Immunofluorescence (1:50-1:100)