

36-3606: Anti-CD40 / TNFRSF5 / CD40L-Receptor Monoclonal Antibody(Clone: C40/1605)

Clonality :	Monoclonal
Clone Name :	C40/1605
Application :	IF,FACS,IHC
Reactivity :	Human
Gene :	CD40
Gene ID :	958
Uniprot ID :	P25942
Alternative Name :	B-cell surface antigen CD40; Bp50; CD40; CD40L receptor; GP39; HIGM1; IGM; IMD3; p50; TBAM; TNF receptor superfamily member 5; TNFRSF5; TRAP
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Recombinant full-length human CD40 protein

Description

CD40 is a receptor on antigen-presenting cells of the immune system and is essential for mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin class switching, memory B cell development, and germinal center formation. AT-hook transcription factor AKNA is reported to coordinately regulate the expression of this receptor and its ligand, which may be important for homotypic cell interactions. Adaptor protein TNFR2 interacts with this receptor and serves as a mediator of the signal transduction. The interaction of this receptor and its ligand is found to be necessary for amyloid-beta-induced microglial activation, and thus is thought to be an early event in Alzheimer disease pathogenesis. CD40 is expressed on B-lymphocytes, follicular dendritic cells, bone marrow-derived dendritic cells, thymic epithelium, and interdigitating cells in the T-cell zones of secondary lymphoid organs.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Immunofluorescence (1-2ug/ml); Flow Cytometry (1-2ug/million cells); ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),




Fig. 1: Formalin-fixed, paraffin-embedded human Hodgkin's Lymphoma stained with CD40 Mouse Monoclonal Antibody (C40/1605).




Fig. 2: Formalin-fixed, paraffin-embedded human Spleen stained with CD40 Mouse Monoclonal Antibody (C40/1605).




Fig. 3: Formalin-fixed, paraffin-embedded human Tonsil stained with CD40 Mouse Monoclonal Antibody (C40/1605).




Fig. 4: SDS-PAGE Analysis Purified CD40 Mouse Monoclonal Antibody (C40/1605). Confirmation of Integrity and Purity of Antibody.




Fig. 5: Flow Cytometric Analysis of U2OS cells using CD40 Mouse Monoclonal Antibody (C40/1605) followed by goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).