

36-3631: Anti-CD47 / IAP (Integrin Associated Protein) Monoclonal Antibody(Clone: CD47/3019)

Clonality :	Monoclonal
Clone Name :	CD47/3019
Application :	ELISA,FACS,IF,IHC
Reactivity :	Human
Gene :	CD47
Gene ID :	961
Uniprot ID :	Q08722
Alternative Name :	Antigenic Surface Determinant Protein OA3; IAP; Integrin Associated Protein; Integrin Associated Signal Transducer; MER6; OA3; Protein MER6; Rh Related Antigen
Isotype :	Mouse IgG1, kappa
Immunogen Information :	A recombinant fragment (around aa 18-135) of human CD47 protein (exact sequence is proprietary)

Description

This antibody reacts with Ig domain of CD47 protein. CD47, originally named integrin-associated protein (IAP), is a 50kDa protein containing five membrane-spanning sequences and a short cytoplasmic tail. CD47 plays a role in both cell adhesion by acting as an adhesion receptor for THBS1 on platelets, and in the modulation of integrins. It is important in memory formation and synaptic plasticity in the hippocampus. CD47 may play a role in membrane transport and/or integrin dependent signal transduction.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

ELISA (For coating, order without BSA);,Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),




Fig. 1: Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with CD47 Mouse Monoclonal Antibody (IAP/3019).




Fig. 2: Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with CD47 Mouse Monoclonal Antibody (IAP/3019).




Fig. 3: SDS-PAGE Analysis Purified CD47 Mouse Monoclonal Antibody (IAP/3019). Confirmation of Integrity and Purity of Antibody.




Fig. 4: Flow Cytometric Analysis of MCF-7 cells using CD47 Mouse Monoclonal Antibody (IAP/3019) followed by goat anti-Mouse IgG-CF488 (Blue); Isotype control (Red).




Fig. 5: Immunofluorescence staining of MCF-7 cells using CD47 Mouse Monoclonal Antibody (IAP/3019) followed by goat anti-Mouse IgG conjugated to CF488 (green). Nuclei are stained with Reddot.




Fig. 6: Analysis of Protein Array containing more than 19,000 full-length human proteins using CD47 Mouse Monoclonal Antibody (CD47/3019) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.