

36-3669: Anti-Cdk1 / p34cdc2 Serine-Threonine Kinase Monoclonal Antibody(Clone: A17.1.1)

Clonality :	Monoclonal
Clone Name :	A17.1.1
Application :	IF,IP,WB,IHC
Reactivity :	Human, Mouse, Rat
Gene :	CDC2
Gene ID :	983
Uniprot ID :	P06493
Alternative Name :	Cdc2, CDC28A, CDK1, CDKN1, CELL CYCLE CONTROLLER CDC2, Cell division control protein 2 homolog, Cell division cycle 2 G1 to S and G2 to M, Cell division protein kinase 1, Cyclin-dependent kinase 1, p34 Protein Kinase, P34CDC2
Isotype :	Mouse IgG2a, kappa
Immunogen Information :	C-Terminal 2/3rds of Xenopus cdc2 expressed in E.coli

Description

Recognizes a 34kDa protein (cdk1), identified as p34cdc2 (a catalytic subunit of Maturation Promoting Factor). Its epitope maps near the C-terminus of the protein and its core is thought to be LGTPNNEV (aa220-227 in murine cdc2). It shows no cross reaction with cdk2 p32. It supports the kinase activity. p34cdc2 plays a crucial role during cell division and is most active during mitosis. It is a serine/threonine kinase, which is activated by cyclin, by dephosphorylation of tyrosine residues. p34cdc2 is inactivated by a tyrosine kinase. This MAb reportedly inhibits the activation of p34cdc2 kinase by cyclins.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Inhibits activation of p34cdc2 kinase by cyclins; Immunofluorescence (1-2ug/ml); Immunoprecipitation; Kinase Assay; Western Blot; Immunohistochemistry (Formalin-fixed) (2-4ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Tonsil stained with Cdk1 Mouse Monoclonal Antibody (A17.1.1).




Fig. 2: Western Blot Analysis of human HeLa cell lysate using Cdk1 Mouse Monoclonal Antibody (A17.1.1).




Fig. 3: SDS-PAGE Analysis Purified Cdk1 Mouse Monoclonal Antibody (A17.1.1). Confirmation of Integrity and Purity of Antibody.