

## 36-3712: Anti-HA Tag Monoclonal Antibody(Clone: 16.43)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	16.43
<b>Application :</b>	ELISA,IF,WB,IP,FACS,IHC
<b>Reactivity :</b>	Human
<b>Alternative Name :</b>	HA epitope tag; HA1; HA2; hemagglutinin; Hemagglutinin HA1 chain; Hemagglutinin HA2 chain
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Semi-Purified mitochondrial preparation

### Description

Human influenza hemagglutinin (HA) is a surface glycoprotein required for the infectivity of the human virus. The HA tag is derived from the HA molecule corresponding to amino acids 98-106 has been extensively used as a general epitope tag in expression vectors. Many recombinant proteins have been engineered to express the HA tag, which does not appear to interfere with the bioactivity or the biodistribution of the recombinant protein. This tag facilitates the detection, isolation, and purification of the proteins.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

ELISA (For coating, order antibody without BSA); Immunofluorescence (1-2ug/ml); Western Blot (1-2ug/ml); Immunoprecipitation (1-2ug per 100-500ug of total protein (1ml of cell lysate)); Flow Cytometry (1-2ug/million cells); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 &degC followed by cooling at RT for 20 minutes),