

36-3714: Anti-Cig2 Monoclonal Antibody(Clone: 3A11/5)

Clonality :	Monoclonal
Clone Name :	3A11/5
Application :	ELISA,WB
Alternative Name :	B type S phase cyclin antibody
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Bacterial <i>S. pombe</i> Cig2 Purified protein.

Description

Cig2 is a cyclin that functions as the catalytic subunit of cyclin-dependent kinases (Cdks) in the fission yeast *Schizosaccharomyces pombe*. It is a B-type, S phase cyclin that is required for both the G1/S and G2/M cell cycle transitions and is expressed in a sharp spike that peaks during the G1/S period. Cig2 binds to Cdc2 (Cdk), and the resulting Cdc2/Cig2 complex controls the G1/S transition of the cell cycle. Disruption of Cig2 delays the onset of mitosis. The expression of the Cig2 gene is dependent on Mlu1-binding factor (MBF), and the protein is destroyed during anaphase by the APC/cyclosome (APC/C) and Skp1/Cullin-1/F-box (SCF), thereby ensuring the spike expression pattern of Cig2. SCF regulates Cig2 levels in a dual manner, transcriptionally and post-translationally, while APC/C only destroys the protein.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

ELISA (For coating, order antibody without BSA);Western Blot (1-2ug/ml);