

36-3718: Anti-Myofibroblast Marker Monoclonal Antibody(Clone: PR 2D3)

Clonality :	Monoclonal
Clone Name :	PR 2D3
Application :	WB,IHC
Reactivity :	Human
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Crude homogenate of normal human colorectal mucosa

Description

Myofibroblasts are a unique group of smooth muscle-like fibroblasts that play an important role in oncogenesis, inflammation, repair, wound contraction and fibrosis. Like smooth muscle (SM) cells, myofibroblasts contain microfilament bundles and express α -SM Actin, the Actin isoform that is present in myoepithelial cells and SM cells and especially abundant in vascular SM cells. Myofibroblasts secrete inflammatory and anti-inflammatory cytokines, chemokines, growth factors and lipid and gaseous inflammatory mediators, as well as extracellular matrix proteins and proteases in most organs and tissues. Besides being temporarily present following tissue injuries and fibrocontractive diseases, myofibroblasts are also present under normal conditions in regions such as the skin, pulmonary septa and periodontal ligaments. Stem cell factor and platelet-derived growth factor (PDGF) are two secreted proteins responsible for differentiating myofibroblasts from embryological stem cells. PR 2D3 reacts with a cell membrane component of cells in the pericypt sheath; with smooth muscle cells and myofibroblasts. This antibody is considered to be the gold standard for the identification of myofibroblasts.

Product Info

Amount :	20 μ g / 100 μ g
Content :	200 μ g/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Western Blot (1-2ug/ml) Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: SDS-PAGE Analysis Purified Myofibroblast Marker Mouse Monoclonal Antibody (PR 2D3). Confirmation of Purity and Integrity of Antibody.




Fig. 2: Western Blot Analysis of A431, MCF-7 & HeLa cell lysates using Myofibroblast Marker Mouse Monoclonal Antibody (PR 2D3).