

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 36-3751: Anti-Campylobacter jejuni Monoclonal Antibody(Clone: 2E-10)

Clone Name: Monoclonal
Clone Name: 2 E-10
Application: ELISA,IF

Alternative Name: Campylobacter jejuni Isotype: Mouse IgM, kappa

Immunogen Information: Total sonicate of Campylobacter jejuni

## **Description**

This MAb is specific for a non-flagellar antigen ofboth C. jejuni Types 1 and 2. It reacts with a soluble excreted antigen in EIA. This determinant is unaffected by frozen storage of specimens, unlike antibodies to flagellar antigens which require fresh cultured organisms. Campylobacter jejuni is a Gram-negative, microaerophilic, slender, flagellate, spiral bacterium. It is the major reported cause of bacterial foodborne infection in the United States and is also associated with Guillian-Barre syndrome. Campylobacteriosis is an infectious disease caused by bacteria of the genus Campylobacter. Most people who become ill with campylobacteriosis get diarrhea, cramping, abdominal pain, and fever within 2 to 5 days after exposure to the organism. The diarrhea may be bloody and can be accompanied by nausea and vomiting. The illness typically lasts 1 week. Some persons who are infected with Campylobacter don't have any symptoms at all. In persons with compromised immune systems, Campylobacter occasionally spreads to the bloodstream and causes a serious life threatening infection. The Campylobacter organism is actually a group of spiral shaped bacteria that can cause disease in humans and animals. Most human illness is caused by one species, called Campylobacter jejuni, but 1% of human Campylobacter cases are caused by other species.

## **Product Info**

**Amount :** 20 μg / 100 μg

Content: 200 µg/ml of Ab Purified from Bioreactor Concentrate. Prepared in 10mM PBS with 0.05% BSA &

0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage condition:

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous.

## **Application Note**

ELISA (For coating use Ab at 1-5ug/ml, order Ab without BSA);Immunofluorescence (1-2ug/ml);