

## 36-3772: Anti-Cytokeratin, Multi (Epithelial Marker) Monoclonal Antibody(Clone: rKRT/457)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	rKRT/457
<b>Application :</b>	IHC
<b>Reactivity :</b>	Human, Rat, Mouse
<b>Gene ID :</b>	3851; 3852; 3853; 3854; 286887; 3856; 3858; 3860; 3875
<b>Uniprot ID :</b>	P02538 ; P04259 ; P13647 ; P19013 ; P48668
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Keratin-enriched preparation from cultured human epithelial cells

### Description

Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) subfamilies. This antibody recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, including 59kDa (CK4); 58kDa (CK5); 56kDa (CK6); 52kDa (CK8); 56.5kDa (CK10); 53kDa (CK13) and 45kDa (CK18). This is a broad-spectrum antibody, which has been reported to differentiate epithelial tumors from non-epithelial tumors. Many studies have shown the usefulness of keratins as markers in cancer research and tumor diagnosis.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Skin stained with Multi-Cytokeratin Mouse Recombinant Monoclonal Antibody (rKRT/457).




Fig. 2: SDS-PAGE Analysis Purified Multi-Cytokeratin Mouse Recombinant Monoclonal Antibody (rKRT/457).