

## 36-3786: Anti-Epstein-Barr Virus (LMP-1) Monoclonal Antibody(Clone: CS4)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	CS4
<b>Application :</b>	IF,WB,IHC
<b>Alternative Name :</b>	EBNA1; EBV nuclear antigen 1; Epstein Barr nuclear antigen 1; Human Herpesvirus 4 (HHV4)
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Recombinant fusion protein containing the sequence of bacterial beta-galactosidase and the carboxyl half of EBV-encoded LMP

### Description

This antibody is a mixture of four different monoclonal antibodies. This antibody is specific to 60kDa latent membrane protein (LMP-1) encoded by the BNLF1 gene of the EBV. Each clone reacts with different epitopes on the hydrophilic C-terminus of the cytoplasmic domain of LMP-1. This antibody stains strongly with EBV-positive lymphoblastoid cell lines and EBV infected B cell immunoblasts in infectious mononucleosis. EBV, also designated human herpesvirus 4 (HHV-4), is a member of the herpesvirus family and is one of the most common human viruses. EBV infects B cells and, though often asymptomatic, it can cause infectious mononucleosis, a disease characterized by fatigue, fever, sore throat and muscle soreness.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Immunofluorescence (1-2ug/ml); Western Blot (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes);




Fig. 1: SDS-PAGE Analysis Purified EBV Mouse Monoclonal Antibody (CS4). Confirmation of Integrity and Purity of Antibody.




Fig. 2: Formalin-fixed, paraffin-embedded human Hodgkin's Lymphoma stained with EBV Mouse Monoclonal Antibody (CS-4).