

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-3821: Anti-CD40, Mouse Monoclonal Antibody(Clone: FGK45.5)

Clonality: Monoclonal Clone Name: FGK45.5

Application: Functional Assay, FACS, IF

Reactivity: Mouse
Gene ID: 21939
Uniprot ID: P27512

Alternative Name:

B cell surface antigen CD40, CD40 type II isoform, p50, Bp50, CD40L receptor, CDW40, TRAP,

IMD3, IGM, Tnfrsf5, HIGM1, GP39, T-BAM

Isotype: Rat IgG2a, kappa

Immunogen Information: Purified recombinant Mouse CD40 fusion protein

Description

Upon binding to its ligand CD154, CD40 acts as a costimulatory molecule for the activation of B cells, dendritic cells, monocytes, and other APCs. CD40 plays roles in B cell activation, differentiation, proliferation and Ig isotype switching as well as dendritic cell maturation. FGK4.5 can be used for blocking of CD40/CD154 interaction and forin vitroandin vivoactivation of CD40 expressing APCs. This MAb is shown to indirectly activate natural killer (NK) cells, producing significant antitumor and anti-metastatic effects. Effective in boosting immune responses against infectious agents and can potentially be used to treat chronic autoimmune inflammatory processes.

Product Info

Amount: 20 μg / 100 μg

Content: 200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS

with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage condition : Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous.

Application Note

Functional Studies (Order Antibody without BSA & Azide); Flow Cytometry (0.5-1ug/million cells in 0.1ml); Immunofluorescence (1-2ug/ml);