

### 35-1883: SARS-CoV Spike Antibody, Rabbit pAb (Cross react with RBD domain) (Discontinued)

<b>Clonality :</b>	Polyclonal
<b>Application :</b>	ELISA
<b>Alternative Name :</b>	Anti-coronavirus s1 Antibody; Anti-coronavirus s2 Antibody; Anti-coronavirus spike Antibody; Anti-cov spike Antibody; Anti-ncov RBD Antibody; Anti-ncov s1 Antibody; Anti-ncov s2 Antibody; Anti-ncov spike Antibody; Anti-novel coronavirus RBD Antibody; Anti-novel coronavirus s1 Antibody; Anti-novel coronavirus s2 Antibody; Anti-novel coronavirus spike Antibody; Anti-RBD Antibody; Anti-S1 Antibody; Anti-s2 Antibody; Anti-Spike RBD Antibody
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Recombinant SARS-CoV Spike S1 Subunit Protein

#### Description

The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors on the host cell. Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2; DPP4, dipeptidyl peptidase-4; APN, aminopeptidase N; CEACAM, carcinoembryonic antigen-related cell adhesion molecule 1; Sia, sialic acid; O-ac Sia, O-acetylated sialic acid. The spike is essential for both host specificity and viral infectivity. The term 'peplomer' is typically used to refer to a grouping of heterologous proteins on the virus surface that function together. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. It's been reported that 2019-nCoV can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity. The main functions for the Spike protein are summarized as: Mediate receptor binding and membrane fusion; Defines the range of the hosts and specificity of the virus; Main component to bind with the neutralizing antibody; Key target for vaccine design; Can be transmitted between different hosts through gene recombination or mutation of the receptor binding domain (RBD), leading to a higher mortality rate.

#### Product Info

<b>Amount :</b>	50 µl / 100 µl
<b>Purification :</b>	Protein A & Antigen Affinity
<b>Content :</b>	0.2 um filtered solution in PBS
<b>Storage condition :</b>	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Avoid repeated freeze-thaw cycles.

#### Application Note

WB 1:1000-1:5000 ELISA 1:5000-1:10000 Specificity: SARS CoV spike/S Has cross-reactivity in ELISA and WB with SARS-CoV-2 (2019-nCoV) Spike S1 SARS-CoV-2 (2019-nCoV) Spike RBD