

## 32-190025: Human ACE2 Protein His Tag (1-740 aa)

**Application :** Functional Assay  
**Uniprot ID :** Q9BYF1  
**Alternative Name :** SARS Receptor; Angiotensin-converting Enzyme 2; ACEH; Metalloprotease MPROT15; SARS-CoV-2 Receptor,

### Description

**Source:** HEK 293 cells. Angiotensin-converting enzyme 2 (ACE2) is an ectoenzyme (carboxypeptidase) with an extracellular catalytic domain that predominantly localizes at the plasma membrane and is thereby able to hydrolyze circulating peptides. ACE2 has approximately 42% sequence identity with ACE, and its cytoplasmic and transmembrane domains show 48% homology to the protein collectrin that plays a critical role in the amino acid absorption of the kidney. ACE2 converts angiotensin I to angiotensin 1-9, a peptide of unknown function, and angiotensin II to angiotensin 1-7, a vasodilator. ACE2 is involved in the regulation of systemic blood pressure and has direct effects on cardiac functions. It is expressed predominantly in endothelial cells of the lung, gut, heart and kidney. ACE2 together with the protease TMPRSS2 acts as a functional receptor for SARS coronavirus as well as for the new highly pathogenic coronavirus, 2019-nCoV/SARS-CoV-2, which is cause for pneumonia COVID-19. It has been shown that human recombinant soluble ACE2 can significantly block early stages of SARS-CoV-2 infections.

### Product Info

**Amount :** 100 µg  
**Purification :** ≥90% (SDS-PAGE)  
**Content :** 1mg/ml after reconstitution. Reconstitute with 100µl endotoxin-free water.  
Formulation:Lyophilized. Contains PBS.  
**Storage condition :** Short Term Storage +4°C. Long Term Storage -20°C. After opening, prepare aliquots and store at -20°C. Avoid freeze/thaw cycles. For maximum product recovery after thawing, centrifuge the vial before opening the cap. Use/Stability-Stable for at least 6 months after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.  
**Amino Acid :** Extracellular domain of human ACE2 (aa 1-740) is fused at the C-terminus to a His-tag.

### Application Note

Biological Activity: Binds to the Spike protein of the coronavirus SARS-CoV-2. Endotoxin Content: <0.001EU/µg purified protein (LAL test).