

12-4031: Phospho-Btk (Tyr223) (Clone: B4) rabbit mAb FITC conjugate

Clonality :	Monoclonal
Clone Name :	BtkY223-B4
Application :	FACS
Reactivity :	Human, Mouse
Conjugate :	FITC
Format :	Conjugated
Alternative Name :	Bruton tyrosine kinase, Tyrosine-protein kinase BTK, Agammaglobulinemia tyrosine kinase, ATK, AGMX1, B-cell progenitor kinase, BPK
Isotype :	Rabbit IgG1k
Immunogen Information :	A synthetic phospho-peptide corresponding to residues surrounding Tyr223 of human phospho Btk

Description

Btk is a major node in the B-cell receptor signaling pathway, where it regulates B cell maturation, activation, survival, differentiation, and proliferation. Btk is activated by Src family kinases, including Lyn, which phosphorylates Btk at Tyr551. Upon phosphorylation at this site, Btk is recruited to the plasma membrane where autophosphorylation at Tyr223 occurs. The Btk signaling pathway is a major target of small molecule inhibitors for B-cell lymphoma, autoimmune diseases, and non-Hodgkin's lymphomas. These inhibitors either form a covalent bond at Cys481 in the ATP-binding site or serve as reversible inhibitors that bind the SH3 pocket and stabilize inactive Btk.

Product Info

Amount :	10 Tests / 100 Tests
Content :	1X PBS, 0.09% NaN ₃ , 0.2% BSA
Storage condition :	Store at 2-8°C. Avoid repeated freeze and thaw cycles.

Application Note

For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells or 5 μ L per 100 μ L of staining volume. It is recommended that the reagent be titrated for optimal performance for each application.




Fig-1: Flow cytometric analysis of U937 cells unstained and untreated cells as negative control (blue) or untreated (red) or treated with IFNa IL4 and PV (green) Phospho-Btk (Tyr223) antibody, BtkY223-B4 FITC conjugate.