

12-4233: Phospho-Stat4 (Tyr693) (Clone: F6) rabbit mAb PE conjugate

Clonality :	Monoclonal
Clone Name :	Stat4Y693-F6
Application :	FACS
Reactivity :	Human, Mouse
Conjugate :	PE
Format :	Conjugated
Alternative Name :	Signal transducer and activator of transcription 4
Isotype :	Rabbit IgG1k
Immunogen Information :	A synthetic phospho-peptide corresponding to residues surrounding Tyr693 of human phospho Stat4

Description

In response to IL-12 binding, the IL-12 receptor activates the Jak kinases, which phosphorylate tyrosine residues of IL-12RB2. These phosphorylated receptors recruit Stat4 through its SH2 domain, whereupon Stat4 is phosphorylated at Tyr693 in its C-terminal transactivation domain. Phosphorylation promotes Stat4 homodimerization and translocation to the nucleus, where it promotes gene transcription. The N-terminal domain of Stat4 appears to be required for maximal stabilization and for the binding of Stat4 dimers to lower-affinity DNA binding sites. Stat4-deficient mice have demonstrated that this gene is required to both promote Th1 development and inhibit Th2 differentiation due to disabling IL-12 receptor-mediated responses.

Product Info

Amount :	10 Tests / 100 Tests
Content :	1X PBS, 0.09% NaN ₃ , 0.2% BSA
Storage condition :	Store at 2-8°C. Avoid repeated freeze and thaw cycles.

Application Note

For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells or 5 μ L per 100 μ L of staining volume. It is recommended that the reagent be titrated for optimal performance for each application.




Fig-1: Flow cytometric analysis of K562 cells unstained imatinib treated cells negative control (blue) or stained treated with imatinib (red) or treated with IFN α + IL-4 + pervanadate (green) using phospho-Stat4 (Tyr693) antibody Stat4Y693-F6 PE conjugate.