

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-1643: Monoclonal Antibody to Bcl-2 (Apoptosis & Follicular Lymphoma Marker)(Clone: SPM530)

Clonality: Monoclonal Clone Name: SPM530

Application : FACS,IF,WB,IHC

Reactivity: Human
Gene: BCL2
Gene ID: 596
Uniprot ID: P10415
Format: Purified
Alternative Name: BCL2

Isotype: Mouse IgG1, kappa

Immunogen Information: A synthetic peptide, aa41-54 (GAAPAPGIFSSQPG-Cys) of human Bcl-2 protein

Description

This antibody recognizes a protein of 25-26kDa, identified as the bcl-2 α oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of bcl-2 α oncoprotein inhibits the programmed cell death (apoptosis). In most follicular lymphomas, neoplastic germinal centers express high levels of bcl-2 α protein, whereas the normal or hyperplastic germinal centers are negative. Consequently, this antibody is valuable when distinguishing between reactive and neoplastic follicular proliferation in lymph node biopsies. It may also be used in distinguishing between those follicular lymphomas that express bcl-2 protein and the small number in which the neoplastic cells are bcl-2 negative.

Product Info

Amount: 100 μg

Purification: Affinity Chromatography

Content: 100 μg in 500 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly


toxic.

Storage condition : Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Western Blot (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),




Formalin-fixed, paraffin-embedded human non-Hodgkin's Lymphoma stained with Bcl-2 Monoclonal Antibody (SPM530).




9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com



Flow Cytometry of Jurkat cells using Bcl-2 Monoclonal Antibody (SPM530) (red) and isotype control (green).



Western Blot of Bcl-2 in human Skin using Bcl-2 Monoclonal Antibody (SPM530).