

## 12-4297: Phospho-mTOR (Ser2448) (Clone: E11) rabbit mAb FITC Conjugate

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	mTORS2448-E11
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human
<b>Conjugate :</b>	FITC
<b>Alternative Name :</b>	Serine/threonine-protein kinase mTOR, FK506-binding protein 12-rapamycin complex-associated protein 1, FKBP12-rapamycin complex-associated protein, Mammalian target of rapamycin, Mechanistic target of rapamycin, Rapamycin and FKBP12 target 1, Rapamycin target protein 1, FRAP, FRAP1, FRAP2, RAFT1, RAPT1
<b>Isotype :</b>	Rabbit IgG1k
<b>Immunogen Information :</b>	A synthetic phospho-peptide corresponding to residues surrounding Ser2448 of human phospho mTOR

### Description

mTOR, mammalian target of rapamycin, is a Serine/Threonine protein kinase (1-2) that functions as an amino acid and ATP sensor to balance cell growth and nutrient availability (3-4). When sufficient nutrients are available, mTOR transmits a positive signal to p70 S6 kinase and participates in the inactivation of 4E-BP1 (5). mTOR plays a key role in homeostasis and cell growth, and phospho mTOR may be abnormally regulated in tumors. mTOR is a potential target for anti-cancer therapy (6).

### Product Info

<b>Amount :</b>	10 Tests / 100 Tests
<b>Content :</b>	1X PBS, 0.09% NaN <sub>3</sub> , 0.2% BSA
<b>Storage condition :</b>	Store at 2-8°C. Avoid repeated freeze and thaw cycles.

### Application Note

For flow cytometric staining, the suggested use of this reagent is 5  $\mu$ L per million cells or 5  $\mu$ L per 100  $\mu$ L of staining volume. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.




Fig-1: Flow cytometric analysis of A431 cells treated with phosphatase and unstained as negative control (blue) or treated with phosphatase (red) or EGF (green) and stained using Phospho-mTOR (Ser2448) FITC conjugated antibody mTORS2448-E11.