

**32-190044: IL-6 (human):Fc (human) (Recombinant)**

<b>Application :</b>	Functional Assay
<b>Reactivity :</b>	Human
<b>Alternative Name :</b>	Interleukin-6; B Cell Stimulatory Factor 2; BSF-2; CTL Differentiation Factor; CDF; Interferon beta2; IFN-beta2

**Description**

Source :CHO cells

Specific Interleukin 6 (IL-6) is a pleiotropic alpha-helical cytokine that plays important roles in acute phase reactions, inflammation, hematopoiesis, bone metabolism and cancer progression. IL-6 activity is essential for the transition from acute inflammation to either acquired immunity or chronic inflammatory disease. It is secreted by multiple cell types as a 22kDa-28kDa phosphorylated and variably glycosylated molecule. IL-6 induces signaling through a cell surface heterodimeric receptor complex composed of a ligand binding subunit (IL-6R) and a signal transducing subunit (gp130). IL-6 is a key factor for the growth of plasma cells.

**Product Info**

<b>Amount :</b>	6 (human):Fc (human) (Recombinant)(Discontinued) / 50 µg
<b>Purification :</b>	>=98% (SDS-PAGE)
<b>Content :</b>	Reconstitute 50µg vial in 50µl sterile water. Add 1X PBS to the desired protein concentration. Lyophilized from 0.2µm-filtered solution in PBS.
<b>Storage condition :</b>	Short Term Storage +4°C ; Long Term Storage-20°C ; Avoid freeze/thaw cycles. Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.
<b>Amino Acid :</b>	The extracellular domain of human IL-6 (aa 30-212) is fused to the N-terminus of the Fc region of human IgG1.

**Application Note**

Endotoxin Content: <0.06EU/µg protein (LAL test; Lonza).