

32-9326: Recombinant Cynomolgus CD27 molecule/CD27(C-6His)(Discontinued)

Alternative Name : CD27 Antigen; CD27L Receptor; T-Cell Activation Antigen CD27; T14; Tumor Necrosis Factor Receptor Superfamily Member 7; CD27; CD27; TNFRSF7

Description

Source : Human Cells;

CD27 antigen, also known as CD27L receptor, T-cell activation antigen CD27, T14, S152, Tp55, TNFRSF7 and Tumor necrosis factor receptor for superfamily member 7, belongs to the TNF-receptor superfamily. CD27 is a single-pass type I membrane protein and exists as a homodimer form, containing three TNFR-Cys repeats. CD27 transduces signals that lead to the activation of NF-KappaB and MAPK8/JNK. CD27 is involved in regulating B-cell activation and immunoglobulin synthesis, binding to the ligand CD70. TRAF2 and TRAF5 have been shown to mediate the signaling process of CD27. CD27-binding protein (SIVA), which is a proapoptotic protein, can bind to CD27 and is thought to play a key role in the apoptosis. CD27 is required for generation and long-term maintenance of T cell immunity.

Product Info

Amount : 500 µg / 50 µg

Content : Lyophilized from a 0.2 µm filtered solution of 20mMTris,150mM NaCl,pH8.5.

Amino Acid : Recombinant Cynomolgus CD27 molecule is produced by our Mammalian expression system and the target gene encoding Thr21-Ile192 is expressed with a 6His tag at the C-terminus.