

32-9352: Recombinant Cynomolgus Programmed cell death ligand 2/PD-L2(C-Fc)

Alternative Name : Programmed Cell Death 1 Ligand 2; PD-1 Ligand 2; PD-L2; PDCD1 Ligand 2; Programmed Death Ligand 2; Butyrophilin B7-DC; B7-DC; CD273; PDCD1LG2; B7DC; CD273; PDCD1L2; PDL2

Description

Source : Human Cells;

Programmed Cell Death 1 Ligand 2 (PDCD1LG2) is a member of the BTN/MOG family. PDCD1LG2 contains one Ig-like C2-type domain and one Ig-like V-type domain. PDCD1LG2 is highly expressed in the heart, placenta, pancreas, lung and liver; it is weakly expressed in the spleen, lymph nodes, and thymus. PDCD1LG2 is involved in the costimulatory signal, essential for T-cell proliferation and IFNG production in a PDCD1-independent manner. PDCD1LG2 interaction with PDCD1 inhibits T-cell proliferation by blocking cell cycle progression and cytokine production.

Product Info

Amount : 500 µg / 50 µg

Content : Lyophilized from a 0.2 µm filtered solution of PBS,pH7.4.

Amino Acid : Recombinant Cynomolgus monkey Programmed cell death ligand 2 is produced by our Mammalian expression system and the target gene encoding Leu20-Pro219 is expressed with a Fc tag at the C-terminus.