

## 32-9398: Recombinant Human Receptor-Type Tyrosine-Protein Kinase FLT3/FLT3 (C-Fc)

**Alternative Name :** Receptor-Type Tyrosine-Protein Kinase FLT3; FL Cytokine Receptor; Fetal Liver Kinase-2; FLK-2; Fms-Like Tyrosine Kinase 3; FLT-3; Stem Cell Tyrosine Kinase 1; STK-1; CD135; FLT3; FLK2; STK1

### Description

Source : Human Cells;

The Flt-3 (fms-like tyrosine kinase) receptor, also named Flk-2 and Stk-1 is a member of the class III subfamily of receptor tyrosine kinases that also includes KIT, the receptor for SCF and FMS, the receptor for M-CSF. The extracellular region of these receptors contains five immunoglobulin-like domains and the intracellular region contains a split kinase domain. Human Flt-3 cDNA encodes a 993 amino acid (aa) residue type I membrane protein with a 26 aa residue signal peptide, a 515 aa extracellular domain with 10 potential N-linked glycosylation sites, a 21 aa residue transmembrane domain and a 431 aa residue cytoplasmic domain. Flt-3 expression has been detected in various tissues, including placenta, gonads, and tissues of nervous and hematopoietic origin. Among hematopoietic cells, the expression of Flt-3 was found to be restricted to the highly enriched stem/progenitor cell populations. The ligand for Flt-3 (FL) has been identified to be a transmembrane protein with structural homology to M-CSF and SCF. Recombinant soluble Flt-3/Fc chimeric protein has been shown to bind FL with high affinity and is a potent FL antagonist.

### Product Info

**Amount :** 500 µg / 50 µg

**Content :** Supplied as a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH7.2.

**Amino Acid :** Recombinant Human Receptor-Type Tyrosine-Protein Kinase FLT3/FLT3 is produced by our Mammalian expression system and the target gene encoding Asn27-Asn541 is expressed with a Fc tag at the C-terminus.Å