

32-9727: Recombinant Human CD28/TP44 (C-Fc)

Alternative Name : CD28; CD28 antigen; CD28 molecule; T-cell-specific surface glycoprotein CD28; Tp44; TP44

Description

Source : Human Cells;

T-cell-specific surface glycoprotein CD28(CD28) is a single-pass type1 membrane protein which contains one Ig-likeV-type (immunoglobulin-like) domain. It belongs to the immunoglobulin(Ig) superfamily. CD28 is one of the molecules expressed on T cells that provide co-stimulatory signals, which are required for T cell activation.CD28 co-stimulation is necessary for CD4 positive T-cell proliferation and survival, interleukin-2 production, and T-helper type-2 development. Human post-thymic regulatory T cells require CD28 co-stimulation to expand and maintain potent suppressive function in vivo. Apoptosis plays a key role in the age-related decline of CD28 expression and in immunosenescence. CD28 is the receptor for CD80 (B7.1) and CD86 (B7.2). When activated by Toll-like receptor ligands, the CD80 expression is upregulated in antigen presenting cells (APCs). The CD86 expression on antigen presenting cells is constitutive. CD28 is the only B7 receptor constitutively expressed on naive T cells.

Product Info

Amount : 500 µg / 50 µg

Content : Lyophilized from a 0.2 µm filtered solution of PBS,pH7.4.

Amino Acid : Recombinant Human CD28 is produced by our Mammalian expression system and the target gene encoding Asn19-Pro152 is expressed with a Fc tag at the C-terminus.