

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-6403: IL1A Human, His Active

**Application:** Functional Assay

Interleukin-1 alpha, IL-1 alpha, Hematopoietin-1, Interleukin-1 alpha: Interleukin-1a, IL-1a,

Alternative Name: Interleukin-1 alpha, IL1A, interleukin 1 alpha, IL1F1, IL-1A, preinterleukin 1 alpha, pro-interleukin-1-

alpha.

## **Description**

Source: E.coli.

Sterile Filtered colorless solution.

IL-1 alpha is derived fromactivated macrophages. IL1A proteins take part in the inflammatory response, being identified as endogenous pyrogens, and stimulate the release of prostaglandin and collagenase from synovial cells.IL-1 alpha stimulates thymocyte proliferation by inducing il-2 release, b-cell maturation and proliferation, and fibroblast growth factor activity. IL1A Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 197 amino acids (113-271 a.a) and having a molecular mass of 22.4kDa. IL1A is fused to a 38 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

## **Product Info**

Amount:  $2 \mu g / 10 \mu g$ 

**Purification:** Greater than 90% as determined by SDS-PAGE.

Content: IL1A protein solution (1mg/ml) contains 20mM Tris-HCl (pH 7.5) and 10% glycerol.

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods

**Storage condition:** of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or

BSA). Avoid multiple freeze-thaw cycles.

Amino Acid: MRGSHHHHHH GMASMTGGQQ MGRDLYDDDD KDRWGSHMSA PFSFLSNVKY NFMRIIKYEF

ILNDALNQSI IRANDQYLTA AALHNLDEAV KFDMGAYKSS KDDAKITVIL RISKTQLYVT AQDEDQPVLL KEMPEIPKTI TGSETNLLFF WETHGTKNYF TSVAHPNLFI ATKQDYWVCL AGGPPSITDF QILENQA.

## **Application Note**

Measured in a cell proliferation assay using D10.G4.1 mouse helper T cell. The ED50 for this effect is less or equal to 0.04 ng/ml.