

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-6491: MCSF Mouse, Sf9

Application: Functional Assay

Alternative Name: M-Csf, Macrophage colony-stimulating factor 1, CSF-1, MCSF, Csf1, C87615, MCSF, op,

Processedmacrophage colony-stimulating factor 1.

Description

Source: Sf9, Baculovirus cells. Sterile Filtered colorless solution.

Macrophage Colony Stimulating Factor (M CSF) or colony stimulating factor 1, is a cytokine, that promoted differentiation in hematopoietic stem cells to macrophages. Another role of M CSF is to bind to its receptor (colony stimulating factor 1 receptor) and to take part in placenta growth and development.

MCSF Mouse produced in Sf9 Baculovirus cells is a single, glycosylated polypeptide chain containing 164 amino acids (33-187 aa) and having a molecular mass of 19.1 kDa.MCSF is fused to a 9 amino acid His tag at C-terminus and purified by proprietary chromatographic techniques.

Product Info

Amount: 2 μg / 10 μg

Purification : Greater than 90.0% as determined by SDS-PAGE.

Content: The MCSF solution (1mg/ml) contains 10% Glycerol and Phosphate-Buffered Saline (pH 7.4).

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods

Storage condition: of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or

BSA). Avoid multiple freeze-thaw cycles.

Amino Acid: ADPKEVSEHC SHMIGNGHLK VLQQLIDSQM ETSCQIAFEF VDQEQLDDPV CYLKKAFFLVQDIIDETMRF

KDNTPNANAT ERLQELSNNL NSCFTKDYEE QNKACVRTFH ETPLQLLEKIKNFFNETKNL LEKDWNIFTK

NCNNSFAKCS SRDVVTKPHH HHHH

Application Note

Determined by cell proliferation assay using M-NFS-60 mouse myelogenous leukemia lymphoblast cells. ED50 range for this effect is <= 4ng/ml.