

32-6593: VEGF Mouse (121 a.a.), Yeast

Application : Functional Assay

Alternative Name : Vascular endothelial growth factor A, VEGF-A, Vascular permeability factor, VPF, VEGF, MGC70609.

Description

Source: *Saccharomyces cerevisiae*

Sterile Filtered White lyophilized (freeze-dried) powder.

Vascular endothelial growth factor is an important signaling protein involved in both angiogenesis and vasculogenesis. VEGF activity has been mostly studied on cells of the vascular endothelium, although it does have effects on a number of other cell types (e.g. stimulation monocyte/macrophage migration, neurons, cancer cells, kidney epithelial cells). VEGF mediates increased vascular permeability, induces angiogenesis, vasculogenesis and endothelial cell growth, promotes cell migration, and inhibits apoptosis. In vitro, VEGF has been shown to stimulate endothelial cell mitogenesis and cell migration. VEGF is also a vasodilator and increases microvascular permeability and was originally referred to as vascular permeability factor. Elevated levels of VEGF is linked to POEMS syndrome, also known as Crow-Fukase syndrome. Mutations in VEGF have been associated with proliferative and nonproliferative diabetic retinopathy.

Vascular Endothelial Growth Factor (121 a.a.) Mouse Recombinant produced in yeast is a disulfide-linked homodimer consisting of 2x121 amino acid polypeptide chains, having a molecular mass of approximately 20.7kDa each. VEGF (121 a.a.) is purified by proprietary chromatographic techniques.

Product Info

Amount : 100 µg / 250 µg

Purification : Greater than 95.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4.

Content : It is recommended to reconstitute the lyophilized Vascular Endothelial Growth Factor (121 a.a.) in sterile 18M Omega -cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Storage condition : Lyophilized VEGF (121 a.a.) although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Vascular Endothelial Growth Factor (121 a.a.) should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.

Amino Acid : MAPTTEGEQK SHEVIKFM DV YQRSYCRPIE TLVDIFQEYP DEIEYIFKPS CVPLMRCAGC CNDEALECVP TSESNITMQI MRIKPHQSQH IGEMSFLQHS RCECRPKKDR TKPEKCDKPR R.

Application Note

The activity was measured in a cell proliferation assay using HUVEC human umbilical vein endothelial cells and was found to be 1-4 ng/ml.