

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-6598: WISP2 Human

WNT1 Inducible Signaling Pathway Protein 2, Connective Tissue Growth Factor-Related Protein 58, **Alternative Name :** Connective Tissue Growth Factor-Like Protein, CCN Family Member 5, CTGF-L, CT58, CCN5, WISP-2,

CTGFL, WISP2.

## **Description**

Source: Escherichia Coli.

Sterile Filtered White lyophilized (freeze-dried) powder.

WNT1-inducible-signaling pathway protein 2 (WISP2) belongs to the WNT1 inducible signaling pathway (WISP) protein subfamily, which belongs to the connective tissue growth factor (CTGF) family. The CTGF family members are characterized by 4 conserved cysteine-rich domains: insulin-like growth factor-binding domain, von Willebrand factor type C module, thrombospondin domain and C-terminal cystine knot-like (CT) domain. WISP2 protein lacks the CT domain which is implicated in dimerization binding. WISP2 is possibly involved in bone remodeling. WISP2 is expressed in primary osteoblasts and fibroblasts. WISP2 stimulates osteoblast adhesion and inhibits osteocalcin production. WISP2 expression in colon tumors is reduced while the other 2 WISP members are overexpressed in colon tumors. WISP2 may play an imperative role in modulating bone turnover.

WISP2 Human Recombinant produced in E.coli is a single, non-glycosylated polypeptide chain containing a total of 228 amino acids and having a molecular mass of 24.4kDa.

## **Product Info**

Amount:  $5 \mu g / 20 \mu g$ 

**Purification :** Greater than 95.0% as determined by SDS-PAGE.

Lyophilized from a sterile (0.2μm) filtered aqueous solution containing 0.1% Trifluoroacetic Acid

(TFA).

Content: It is recommended to reconstitute the lyophilized WISP-2 in sterile 10mM acetic acid not less

than 100µg/ml, which can then be further diluted to other aqueous solutions.

Lyophilized WISP2 although stable at room temperature for 3 weeks, should be stored

desiccated below -18°C. Upon reconstitution WISP-2 should be stored at 4°C between 2-7 days

**Storage condition :**and for future use below -18 °C. For long term storage it is recommended to add a carrier protein

(0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Amino Acid: MQLCPTPCTC PWPPPRCPLG VPLVLDGCGC CRVCARRLGE PCDQLHVCDA SQGLVCQPGA

GPGGRGALCL LAEDDSSCEV NGRLYREGET FQPHCSIRCR CEDGGFTCVP LCSEDVRLPS WDCPHPRRVE

VLGKCCPEWV CGQGGGLGTQ PLPAQGPQFS GLVSSLPPGV PCPEWSTAWG PCSTTCGLGM

ATRVSNQNRF CRLETQRRLC LSRPCPPSRG RSPQNSAF.