

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-6767: GLUL Human, His Active

Application: Functional Assay

Alternative Name: GLNS, EC 6.3.1.2, EC 4.1.1.15, GLUL, Glutamine Synthetase, GS, Glutamate decarboxylase,

Glutamate--ammonia ligase, PIG43, PIG59.

Description

Source: Escherichia Coli. Sterile filtered colorless solution.

GLUL catalyzes the synthesis of glutamine from glutamate and ammonia. Glutamine is a major Source: of energy and that takes part in cell proliferation, inhibition of apoptosis, and cell signaling. GLUL is expressed during early fetal stages, and has a role in maintaining body pH by removing ammonia from circulation. Mutations in GLUL gene are related with congenital glutamine deficiency.

GLUL Recombinant Human produced in E.Coli is a single, non-glycosylated polypeptide chain containing 393 amino acids (1-373 a.a.) and having a molecular mass of 44.2 kDa. The GLUL is fused to a 20 amino acid His-Tag at N-terminus and purified by proprietary chromatographic techniques.

Product Info

Amount: $2 \mu g / 10 \mu g$

Purification : Greater than 90.0% as determined by SDS-PAGE.

Content: GLUL Human solution containing 20mM Tris-HCl pH-8, 5mM DTT, 0.2M NaCl & 20% glycerol.

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods

Storage condition: of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or

BSA). Avoid multiple freeze-thaw cycles.

Amino Acid: MGSSHHHHHH SSGLVPRGSH MTTSASSHLN KGIKQVYMSL PQGEKVQAMY IWIDGTGEGL

RCKTRTLDSE PKCVEELPEW NFDGSSTLQS EGSNSDMYLV PAAMFRDPFR KDPNKLVLCE VFKYNRRPAE

TNLRHTCKRI MDMVSNQHPW FGMEQEYTLM GTDGHPFGWP SNGFPGPQGP YYCGVGADRA

YGRDIVEAHY RACLYAGVKI AGTNAEVMPA QWEFQIGPCE GISMGDHLWV ARFILHRVCE DFGVIATFDP KPIPGNWNGA GCHTNFSTKA MREENGLKYI EEAIEKLSKR HQYHIRAYDP KGGLDNARRL TGFHETSNIN

DFSAGVANRS ASIRIPRTVG OEKKGYFEDR RPSANCDPFS VTEALIRTCL LNETGDEPFQ YKN.

Application Note

Specific activity is > 2,800 pmol/min/ug, and is defined as the amount of enzyme that convert 1.0 pmole of L-glutamate to L-glutamine per miunte at pH 7.5 at 37C in coupled system with PK/LDH.