

32-6992: PGK1 Human, Active

Application : Functional Assay

Alternative Name : Phosphoglycerate kinase 1, MGC117307, MGC142128, MGC8947, MIG10, PGKA.

Description

Source: E.coli.

Sterile Filtered colorless solution.

Phosphoglycerate kinase 1, also referred to PGK1, is an X-linked enzyme which functions in the glycolytic pathway. The gene encoding the erythrocyte enzyme PGK1 & it seems that PGK-1 plays a role as a polymerase alpha cofactor protein (primer recognition protein) as a glycolytic enzyme role. The PGK1 catalyzes the reversible conversion of 1, 3-diphosphoglycerate to 3-phosphoglycerate during glycolysis, generating one molecule of ATP.

PGK1 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 437 amino acids (1-417a.a.) and having a molecular mass of 46.8kDa. PGK1 is fused to a 20 amino acid His tag at N-Terminus and purified by proprietary chromatographic techniques.

Product Info

Amount : 2 µg / 10 µg

Purification : Greater than 95.0% as determined by SDS-PAGE.

Content : The PGK1 solution (1mg/ml) contains 10% glycerol, 20mM Tris-HCl buffer (pH 8.0) and 1mM DTT. .

Storage condition : Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Amino Acid : MGSSHHHHHH SSGLVPRGSH MSLSNKLTLD KLDVKGKRVV MRVDFNVPMK NNQITNNQRI
KAAVPSIKFC LDNGAKSVVL MSHLGRPDGV PMPDKYSLEP VAVELKSLG KDVFLKDCV GPEVEKACAN
PAAGSVILLE NLRFHVEEEG KGKDASGNKV KAEPKIEAF RASLSKLDGV YVNDAFGTAH RAHSSMVGVN
LPQKAGGFLM KKELNYFAKA LESPFPFLA ILGGAKVADK IQLINMLDK VNEMIIGGGM AFTFLKVLNN
MEIGTSLFDE EGAKIVKDLN SKAEKNGVKI TLPVDFVTAD KFDENAKTGQ ATVASGIPAG WMGLDCGPES
SKKYAEAVTR AKQIVWNGPV GVFEWEAFAR GTKALMDEVV KATSRGCITI IGGGDTATCC AKWNTEDKVS
HVSTGGGASL ELLEGKVLPG VDALSNI

Application Note

Specific activity: > 600unit/mg. One unit will convert 1 umole of 1,3-Bisphosphoglycerate to 3-PGA per minute at pH 8.0 at 37C.