

32-13359: PDCD1LG2 Human, Sf9

Alternative Name : Programmed Cell Death 1 Ligand 2, B7 Dendritic Cell Molecule, Programmed Death Ligand 2, Butyrophilin B7-DC, PDCD1 Ligand 2, PDCD1L2, B7-DC, CD273, PD-L2,B7DC, PDL2, PD-1-Ligand 2, CD273 Antigen, PD-1 Ligand 2, BA574F11.2, Btdc.

Description

Source: Sf9, Baculovirus cells.

Sterile Filtered colorless solution.

Programmed cell death 1 ligand 2 (PDCD1LG2) which takes part in the costimulatory signal, is necessary for T-cell proliferation. PDCD1LG2 is also involved in IFNG production in a PDCD1-independent way. The interaction with PDCD1 inhibits T-cell proliferation by preventing cytokine production and cell cycle progression.

PDCD1LG2 Human Recombinant produced in Sf9 Baculovirus cells is a single, glycosylated polypeptide chain containing 423 amino acids (20-200a.a.) and having a molecular mass of 47.7kDa (Molecular size on SDS-PAGE will appear at approximately 40-57kDa).PDCD1LG2 is expressed with a 239 amino acids hlgG-His tag at C-Terminus and purified by proprietary chromatographic techniques.

Product Info

Amount : 2 µg / 10 µg

Purification : Greater than 95.0% as determined by SDS-PAGE.

Content : PDCD1LG2 protein solution (1mg/ml) contains Phosphate Buffered Saline (pH 7.4) and 10% glycerol.

Storage condition : Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Avoid multiple freeze-thaw cycles.

Amino Acid : ADPLFTVTVP KELYIIIEHGS NVTLECNFDT GSHVNLGAI ASLQKVENDT SPHRERATLL EEQLPLGKAS FHIPQVQVRD EGQYQCIIY GVAWDYKYLK LKVKASYRKI NTHILKVPET DEVELTCQAT GYPLAEVSWP NVSVPANTSH SRTPEGLYQV TSVLRLKPPP GRNFSCVFWN THVRLEPKSC DKHTCPCPC APELLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD GVEVHNAKTK PREEQYNSTY RVVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK GQPREPQVYT LPPSRDELTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTTPVLDL DGSFFLYSKL TVDKSRWQQG NVFSCSV MHE ALHNHYTQKS LSLSPGKHHH HHH.