

## 12-9021: Anti-CD38 antibody(DM30), Rabbit mAb(Discontinued)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	DM30
<b>Application :</b>	ELISA,FACS
<b>Reactivity :</b>	Human
<b>Alternative Name :</b>	T10, cADPr hydrolase 1
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Recombinant human CD38 (Val43-Ile300) produced by using human HEK293 cells

### Description

CD antigen CD38 is also known as ADP-ribosyl cyclase 1, which belongs to the ADP-ribosyl cyclase family. CD38 is expressed at high levels in pancreas, liver, kidney, brain, testis, ovary, placenta, malignant lymphoma and neuroblastoma. CD38 is a multifunctional ectoenzyme that catalyzes the synthesis and hydrolysis of cyclic ADP-ribose (cADPR) from NAD<sup>+</sup> to ADP-ribose. These reaction products are essential for the regulation of intracellular Ca<sup>2+</sup>. The loss of CD38 function is associated with impaired immune responses, metabolic disturbances, and behavioral modifications. The CD38 protein is a marker of cell activation. It has been connected to HIV infection, leukemias, myelomas, solid tumors, type II diabetes mellitus and bone metabolism. CD38 has been used as a prognostic marker in leukemia.

### Product Info

<b>Amount :</b>	100 µg
<b>Purification :</b>	Purified from cell culture supernatant by affinity chromatography
<b>Content :</b>	Preservative: 0.1% Procline 300 Constituents: 50% Glycerol; PBS,pH 7.4; 0.1% BSA Not Sterile
<b>Storage condition :</b>	Store at -20°C for 12 months (Avoid repeated freezing and thawing)

### Application Note

Recommended Dilutions ELISA 1/5000-10000; FACS 1/100




Figure 1. Expi 293 cell line transfected with irrelevant protein (left) and human CD38 (right) were surface stained with Rabbit anti-CD38 monoclonal antibody 1µg/ml (clone: DM30) followed by Alexa 488-conjugated anti-rabbit IgG secondary antibody.




Figure 2. FACS data of serially titrated Rabbit anti-CD38 monoclonal antibody (clone: DM30) on Raji cells. The Y-axis represents the mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.




Figure 3. Affinity ranking of different Rabbit anti-CD38 mAb clones by titration of different concentration onto Raji cells. The Y-axis represents the mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.