# **w** abeomics

## 32-17058: Recombinant human ACE2 protein with C-terminal human Fc tag

Alternative Name : ACE-2, ACEH, ACE2

### Description


#### Expression Host : HEK293

The protein has a predicted molecular mass of 109.7 kDa after removal of the signal peptide.

The protein encoded by this gene belongs to the angiotensin-converting enzyme family of dipeptidyl carboxydipeptidases and has considerable homology to human angiotensin 1 converting enzyme. This secreted protein catalyzes the cleavage of angiotensin 1 into angiotensin 1-9, and angiotensin II into the vasodilator angiotensin 1-7. The organ- and cell-specific expression of this gene suggests that it may play a role in the regulation of cardiovascular and renal function, as well as fertility. In addition, the encoded protein is a functional receptor for the spike glycoprotein of the human coronaviruses SARS and HCoV-NL63.

#### **Product Info**

Amount :	50 μg
Purification :	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Content :	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose is added as protectants before lyophilization.
Storage condition :	Store at -80°C for 12 months (Avoid repeated freezing and thawing)



10

100

Human ACE2, hFc Tagged protein(ng/ml)

1000

10000

Figure 1. Human ACE2 Protein, hFc Tag on SDS-PAGE under reducing condition.

Figure 2. ELISA plate pre-coated by 2  $\mu$ g/ml (100  $\mu$ l/well) S-RBD, mFc-His tagged protein can bind Human ACE2, hFc Tagged protein in a linear range of 0.488-49.83 ng/ml.