

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-17064: Recombinant human PSCA protein with C-terminal human Fc

Alternative Name: PSCA, UNQ 206, PRO 232

## **Description**

Expression Host: HEK293

The protein has a predicted molecular mass of 34.4 kDa after removal of the signal peptide. The apparent molecular mass of PSCA-hFc is approximately 40-57 kDa due to glycosylation.

This gene encodes a glycosylphosphatidylinositol-anchored cell membrane glycoprotein. In addition to being highly expressed in the prostate it is also expressed in the bladder, placenta, colon, kidney, and stomach. This gene is up-regulated in a large proportion of prostate cancers and is also detected in cancers of the bladder and pancreas. This gene includes a polymorphism that results in an upstream start codon in some individuals; this polymorphism is thought to be associated with a risk for certain gastric and bladder cancers. Alternative splicing results in multiple transcript variants.

## **Product Info**

**Amount:** 50 μg

**Purification:** The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

staining.

**Content:** Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose is added as protectants

before lyophilization.

**Storage condition:** Store at -80°C for 12 months (Avoid repeated freezing and thawing)



Figure 1. Human PSCA Protein, hFc Tag on SDS-PAGE under reducing condition.