

## 32-17064: Recombinant human PSCA protein with C-terminal human Fc

**Alternative Name :** PSCA,UNQ206,PRO232

### Description

Expression Host : HEK293

The protein has a predicted molecular mass of 34.4 kDa after removal of the signal peptide. The apparent molecular mass of PSCA-hFc is approximately 40-57 kDa due to glycosylation.

This gene encodes a glycosylphosphatidylinositol-anchored cell membrane glycoprotein. In addition to being highly expressed in the prostate it is also expressed in the bladder, placenta, colon, kidney, and stomach. This gene is up-regulated in a large proportion of prostate cancers and is also detected in cancers of the bladder and pancreas. This gene includes a polymorphism that results in an upstream start codon in some individuals; this polymorphism is thought to be associated with a risk for certain gastric and bladder cancers. Alternative splicing results in multiple transcript variants.

### Product Info

<b>Amount :</b>	50 µg
<b>Purification :</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Content :</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose is added as protectants before lyophilization.
<b>Storage condition :</b>	Store at -80°C for 12 months (Avoid repeated freezing and thawing)




Figure 1. Human PSCA Protein, hFc Tag on SDS-PAGE under reducing condition.