

32-17086: Recombinant human B7-2 protein with C-terminal human Fc tag

Alternative Name : CD86, B7-2, B70, CD28LG2, LAB72, MGC34413

Description

Expression Host : HEK293

The protein has a predicted molecular mass of 51.5 kDa after removal of the signal peptide.

This gene encodes a type I membrane protein that is a member of the immunoglobulin superfamily. This protein is expressed by antigen-presenting cells, and it is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of this protein with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of this protein with cytotoxic T-lymphocyte-associated protein 4 negatively regulates T-cell activation and diminishes the immune response. Alternative splicing results in several transcript variants encoding different isoforms.

Product Info

- Amount :** 50 µg
- Purification :** The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
- Content :** Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose is added as protectants before lyophilization.
- Storage condition :** Store at -80°C for 12 months (Avoid repeated freezing and thawing)




Figure 1. Human B7-2 Protein, hFc Tag on SDS-PAGE under reducing condition.




Figure 2. ELISA plate pre-coated by 2 µg/ml (100 µl/well) Human B7-2, hFc tagged protein can bind Human CTLA4, mFc-His tagged protein in a linear range of 0.048-2.694 ng/ml.




Figure 3. ELISA plate pre-coated by 2 µg/ml (100 µl/well) Human CD28, mFc-His tagged protein can bind Human B7-2, hFc tagged protein in a linear range of 62.5-4000 ng/ml.




Figure 4. HEK293 cell line transfected with irrelevant protein (B) and human CD28 (A) were surface stained with Human B7-2, hFc tagged protein 1 µg/ml followed by Alexa 488-conjugated anti-human IgG secondary antibody.