

32-17097: SARS-CoV-2 (2019-nCoV) S protein RBD, hFc Tag

Alternative Name : S protein RBD, Spike glycoprotein Receptor-binding domain, S glycoprotein RBD, Spike protein RBD, COVID-19

Description

Expression Host : HEK293

The protein has a predicted molecular mass of 51.2 kDa after removal of the signal peptide.

SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) also known as Covid19 (2019 Novel Coronavirus) is a virus that causes illnesses ranging from the common cold to severe diseases. The spike protein is a type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which accounts for recognizing the cell surface receptor, ACE2. S2 contains basic elements needed for the membrane fusion. Recent publications indicate that S1-RBD domain can induce virus neutralizing-antibody and T cell response.

Product Info

- Amount :** 50 µg
- Purification :** The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
- Content :** Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose is added as protectants before lyophilization.
- Storage condition :** Store at -80°C for 12 months (Avoid repeated freezing and thawing)




Figure 1. SARS-CoV-2 (2019-nCoV) S protein RBD, hFc Tag on SDS-PAGE under reducing condition.




Figure 2. ELISA plate pre-coated by 2 µg/ml (100 µl/well) S-RBD, hFc tagged protein can bind Human ACE2, mFc Tagged protein in a linear range of 0.976-39.35 ng/ml.




Figure 3. ELISA plate pre-coated by 2 µg/ml (100 µl/well) Human ACE2, His Tagged protein can bind S-RBD, hFc tagged protein in a linear range of 0.128-80.0 ng/ml.