

10-4089-P: Monoclonal Antibody to CD3 (Clone: OKT3) PE Conjugated

| | |
|--------------------------------|--|
| Clonality : | Monoclonal |
| Clone Name : | OKT3 |
| Application : | FACS |
| Reactivity : | Human |
| Conjugate : | PE |
| Gene : | CD3D |
| Gene ID : | 915 |
| Uniprot ID : | P04234 |
| Format : | Purified |
| Alternative Name : | CD3D,T3D |
| Isotype : | Mouse IgG2a Kappa |
| Immunogen Information : | Human PBMCs were used as immunogen for the CD3 antibody. |

Description

CD3D is one of the several invariant chains that the variable chains of TCR: Alpha beta or Gamma delta TCR contains, and which support surface TCR expression and transduce the TCR signal. CD3D, along with four other CD3 subunits, binds either TCR alpha/beta or TCR gamma/delta to form the TCR/CD3 complex on the surface of T-cells. Defects in this gene are a cause of severe combined immunodeficiency autosomal recessive T-cell-negative/B-cell-positive/NK-cell-positive (SCIDBNK).

Product Info

| | |
|----------------------------|---|
| Amount : | 25 Test / 100 Test |
| Purification : | Protein G Chromatography |
| Content : | 0.04 mg/ml in PBS with 0.05% Azide. |
| Storage condition : | Store the antibody at 4°C, stable for 6 months. |

Application Note

FACS analysis: 0.1- 0.5 $\mu\text{g}/10^6$ cells




Fig-1: Cell surface flow analysis of CD3 in PBMC using 0.1 $\mu\text{g}/10^6$ cells. Green represents isotype control; red represents anti-hCD3 antibody conjugated with PE (10-4089PE).