

## 30-2486: Anti-Human TCR gamma/delta PE-DyLight® 594 (Clone : B1)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	B1
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human
<b>Gene :</b>	PLCG1
<b>Gene ID :</b>	5335
<b>Alternative Name :</b>	TCRG/D
<b>Isotype :</b>	Mouse IgG1 kappa

### Description

The antigen-specific T cell receptor (TCR) is composed of either alpha and beta subunit, or gamma and delta subunit. Majority of T cells present in the blood, lymph and secondary lymphoid organs express TCR alpha/beta heterodimers, whereas the T cells expressing TCR gamma/delta heterodimers are localized mainly in epithelial tissues and at the sites of infection. The subunits of TCR heterodimers are covalently bonded and in the endoplasmic reticulum they associate with CD3 subunits to form functional TCR-CD3 complex. Lack of expression of any of the chains is sufficient to stop cell surface expression.

**Specificity :** The mouse monoclonal antibody B1 (also known as B1.1) recognizes an extracellular epitope of TCR gamma/delta, the subtype of T cell receptor expressed mainly in epithelial tissues and at the sites of infection.

### Product Info

<b>Amount :</b>	100 tests
<b>Purification :</b>	The purified antibody is conjugated with tandem dye PE-DyLight <sup>Å</sup> ® 594 under optimum conditions. The conjugate is purified by size-exclusion chromatography.
<b>Content :</b>	Formulation : Stabilizing phosphate buffered saline (PBS) solution containing 15 mM sodium azide
<b>Storage condition :</b>	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.

### Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 4 Åµl reagent / 100 Åµl of whole blood or 10<sup>5</sup> cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human TCR gamma/delta (B1) PE-DyLight® 594 antibody




Figure 2: Flow cytometry multicolor surface staining of human lymphocytes stained using anti-human TCR gamma/delta (B1) PE-DyLight® 594 antibody (4 µl reagent / 100 µl of peripheral whole blood) and anti-human CD3 (UCHT1) APC antibody




Figure 3: Separation of human TCR gamma/delta positive T cells (red-filled) from TCR gamma/delta negative CD3 negative lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human TCR gamma/delta (B1) PE-DyLight® 594 antibody