∗ abeomics

30-2576: Anti-Human CD35 APC (Clone : E11)

Clonality :	Monoclonal
Clone Name :	E11
Application :	FACS
Reactivity :	Human
Conjugate :	APC
Gene :	CR1
Gene ID :	1378
Alternative Name :	CR1, KN, C3BR, C4BR,complement C3b/C4b receptor 1 (Knops blood group)
Isotype :	Mouse IgG1
Immunogen Information : Acute monocytic leukemia cells and normal blood monocytes	

Description

Due due to luife

CD35 (complement receptor 1, CR1) is a monomeric multiple modular cell surface glycoprotein which serves as receptor for C3b and C4b, the most important components of the complement system leading to clearance of foreign macromolecules. It is expressed mainly on the surface of granulocytes, monocytes, erythrocytes, B cells and folicular dendritic cells. Besides its role in complement cascade, CD35 is involved in blocking BCR-induced proliferation and the differentiation of B cells to plasmablasts and their Ig production.

Specificity : The mouse monoclonal antibody E11 recognizes an extracellular epitope of CD35 (CR1), a type I transmembrane glycoprotein expressed on granulocytes, monocytes, B cells, folicular dendritic cells, erythrocytes, NK and T cell subsets, as well as e.g. on glomerulal podocytes.

Product Into	
Amount :	100 tests
Purification :	The purified antibody is conjugated with allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography.
Content :	Formulation : Stabilizing phosphate buffered saline (PBS) solution containing 15 mM sodium azide
Storage condition :	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 10 \tilde{A} $\hat{A}\mu$ reagent / 100 \tilde{A} $\hat{A}\mu$ of whole blood or 10⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.




Figure 1 : Flow cytometry analysis (surface staining) of CD35 in human peripheral blood with anti-CD35 (E11) APC.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.