

### 30-2615: Anti-Human CD24 PE-DyLight® 594 (Clone : SN3) (Discontinued)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	SN3
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human
<b>Gene :</b>	CD24
<b>Gene ID :</b>	100133941
<b>Alternative Name :</b>	HSA, nectadorin, SCLC4,CD24 molecule
<b>Isotype :</b>	Mouse IgG1
<b>Immunogen Information :</b>	Glycoproteins purified from human NALM-1 cell line.

#### Description

CD24, also known as heat-stable antigen (HSA) or nectadorin, is a small mucin-like GPI-anchored extracellular membrane glycoprotein expressed on several cell types, including B cells. When B cells are activated and induced to further maturation, however, CD24 begins to disappear. CD24 seems to act as a gate-keeper for lipid rafts, thereby regulating the activity of integrins and other proteins such as the chemokine receptor CXCR4; it is also a ligand for P-selectin. CD24 triggering induces apoptosis of B cell precursors but not in mature resting B cells, where it instead inhibits their ability to proliferate in response to activation. CD24 expression is associated with invasiveness and poorer prognosis of carcinomas and is a marker of exosomes secreted into urine and amniotic fluid.

**Specificity :** The antibody SN3 reacts with CD24, a 35-45 kDa heavily glycosylated cell surface antigen. CD24 is expressed by granulocytes, B lymphocytes and by some activated T cells and T cell malignancies. It is not expressed on human thymocytes.

#### Product Info

<b>Amount :</b>	100 tests
<b>Purification :</b>	The purified antibody is conjugated with tandem dye PE-DyLight <sup>Å</sup> 594 under optimum conditions. The conjugate is purified by size-exclusion chromatography.
<b>Content :</b>	Formulation : Stabilizing phosphate buffered saline (PBS) solution containing 15 mM sodium azide
<b>Storage condition :</b>	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.

#### Application Note

**Flow cytometry:** The reagent is designed for analysis of human blood cells using 4 Åµl reagent / 100 Åµl of whole blood or 10<sup>6</sup> cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.




Figure 1 : Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD24 (SN3) PE-DyLight® 594 antibody (4 µl reagent / 100 µl of peripheral whole blood).




Figure 2 : Separation of human CD24 positive lymphocytes (red-filled) from monocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD24 (SN3) PE-DyLight® 594 antibody (4 µl reagent / 100 µl of peripheral whole blood).