

### 30-2663: Anti-Human CD109 PE (Clone : W7C5)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	W7C5
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human
<b>Conjugate :</b>	PE
<b>Gene :</b>	CD109
<b>Gene ID :</b>	135228
<b>Alternative Name :</b>	CPAMD7, p180, r150, FLJ38569, GPA,CD109 molecule
<b>Isotype :</b>	Mouse IgG1
<b>Immunogen Information :</b>	WERI-RB-1 retinoblastoma cell line

#### Description

CD109, also known as the Gov platelet alloantigen, is a GPI-anchored glycoprotein which localizes to the surface of platelets, activated T-cells, and endothelial cells, as well as of various hematopoietic cells and T cell lines. The protein binds to and negatively regulates signaling by transforming growth factor beta (TGF-beta). Multiple transcript variants encoding different isoforms have been found for this gene. The Gov antigen system is involved in platelet transfusion reaction, posttransfusion purpura and in neonatal alloimmune thrombocytopenia.

Specificity : The mouse monoclonal antibody W7C5 recognizes CD109, an approximately 165 kDa GPI-anchored extracellular protein expressed mainly on various hematopoietic cells, activated T lymphoblasts, activated platelets, and endothelial cells.

#### Product Info

<b>Amount :</b>	100 tests
<b>Purification :</b>	The purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography.
<b>Content :</b>	Formulation : Stabilizing phosphate buffered saline (PBS) solution containing 15 mM sodium azide
<b>Storage condition :</b>	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.

#### Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 10  $\mu\text{l}$  reagent / 100  $\mu\text{l}$  of whole blood or 10<sup>6</sup> cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.




Figure 1 : Flow cytometry surface staining pattern of human PHA stimulated peripheral blood mononuclear cells stained using anti-human CD109 (W7C5) PE antibody (10  $\mu\text{l}$  reagent per million cells in 100  $\mu\text{l}$  of cell suspension).




Figure 2 : Separation of human CD109 positive cells (red-filled) from CD109 negative cells (black-dashed) in flow cytometry analysis (surface staining) of human PHA stimulated peripheral blood mononuclear cells stained using anti-human CD109 (W7C5) PE antibody (10  $\mu$ l reagent per milion cells in 100  $\mu$ l of cell suspension).