

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-2664: Anti-Mouse CD106 Azide Free Antibody (Clone: 429 (MVCAM.A))

Clonality: Monoclonal
Clone Name: 429 (MVCAM.A)

Application : FACS , IP, IHC(F), Functional Assay

Reactivity: Mouse
Gene: Vcam1
Gene ID: 22329
Format: Azide Free

Alternative Name: Vcam-1,vascular cell adhesion molecule 1

Isotype: Rat IgG2a kappa

Immunogen Information: Murine preadipose cell line PA6

Description

CD106 / VCAM-1 (vascular cell adhesion molecule-1) is an Ig-like cell surface adhesion molecule binding VLA-4 integrin. VCAM-1 is a potent T cell costimulatory molecule taking part in their positive selection and survival, as well as in adhesion, transendothelial migration and activation of peripheral T cells. VCAM-1 is also involved in endothelial cell-cell contacts. Whereas VCAM-1 normally mediates leukocyte extravasion to sites of tissue inflammation, tumour cells can use overexpressed VCAM-1 to escape T cell immunity.

Specificity: The rat monoclonal antibody 429 (also known as MVCAM.A) recognizes an extracellular epitope of murine CD106, a 100-110 kDa type I membrane protein of the immunoglobulin superfamily, a crucial mediator of leukocyte adhesion, and a costimulation molecule.

Product Info

Amount: 0.1 mg

Purification : Purified by protein-A affinity chromatography

Content: 1 mg/m

Formulation : Azide free phosphate buffered saline (PBS) solution, 0.2 µm filter sterilized

Storage condition : Store at 2-8°C. Do not freeze.




Figure 1 : Flow cytometry surface staining pattern of murine bone marrow cell suspension stained using anti-mouse CD160 (429) purified antibody (azide free, concentration in sample 0,19 μ g/ml) DAR APC.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2 : Separation of murine CD106 positive cells (red-filled) from CD106 negative cells (black-dashed) in flow cytometry analysis (surface staining) of murine bone marrow cell suspension stained using anti-mouse CD160 (429) purified antibody (azide free, concentration in sample 0,19 μ g/ml) DAR APC.