

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

10-10046: Monoclonal Antibody to SARS-CoV-2 nucleocapsid (Clone: ABM5H9.1A2)

Clonality: Monoclonal Clone Name: ABM5H9.1A2

Application: WB **Gene:** N

Gene ID: 43740575
Uniprot ID: PODTC9
Format: Purified

Isotype: Mouse IgG1, Lambda

Immunogen Information: Full length recombinant SARS-CoV-2 nucleocapsid Protein was used as the immunogen for this

antibody.

Description

The structural nucleocapsid (N) protein of nCoV/SARS-CoV-2/COVID-19 is a predicted 46 kDa phosphoprotein having 419 amino acid residues. A short Serine rich stretch and a recognized nuclear localization signal are the unique features of the nucleocapsid protein of nCoV/SARS-CoV-2/COVID-19, which seems to have a little homology with the proteins of other members of this large corona virus family. These features also suggest the involvement of nucleocapsid protein of nCoV/SARS-CoV-2/COVID-19 in different stages of viral lifecycle. The protein has multifaceted activities including packing of the nCoV/SARS-CoV-2/COVID-19 viral genome into a helical ribonucleocapsid (RNP) and playing an important role in viral self-assembly causing nCoV/SARS-CoV-2/COVID-19. The nucleocapsid protein of nCoV/SARS-CoV-2/COVID-19 also forms dimer in the cell by self-association with the help of interactive C terminal domain.

Product Info

Amount : $25 \mu g / 100 \mu g$

Purification: Protein G Chromatography

Content: 25 μg in 50 μl/100 μg in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium

azide is highly toxic.

Storage condition : Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Recommended dilutions: WB: 0.1-1 μg/ml. However, this need to be optimized based on the research applications.




Figure-1: Western Blot analysis of SARS-CoV-2 Nucleocapsid Antibody: Anti- SARS-CoV-2 Nucleocapsid Antibody (Clone: ABM5H9.1A2) was used at 0.1 μg/ml on (1) Nucleocapsid Recombinant protein and at 1 μg/ml on (2) RBD Recombinant protein.