

## 12-8052: Anti-Human CD52 (Alemtuzumab) - Fc Muted™

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	Campath-1H
<b>Application :</b>	ELISA
<b>Alternative Name :</b>	HE5; CDW52; EDDM5 CDW52; Cambridge pathology 1 antigen
<b>Isotype :</b>	Human IgG1k
<b>Immunogen Information :</b>	Human lymphocytes.

### Description

Expression Host : HEK-293

Pathogen Testing : To protect mouse colonies from infection by pathogens and to assure that experimental preclinical data is not affected by such pathogens, all of this recombinant biosimilar antibodies are tested and guaranteed to be negative for all pathogens in the IDEXX IMPACT I Mouse Profile.

This non-therapeutic biosimilar antibody uses the same variable region sequence as the therapeutic antibody Alemtuzumab. Clone Campath-1H recognizes human CD52. This product is for research use only.

Clone Campath-1H is a monoclonal antibody that specifically binds to CD52, a protein present on the surface of mature lymphocytes. However, this protein is not present on the stem cells that generated these lymphocytes. Alemtuzumab targets and destroys mature lymphocytes containing CD-52, and is used to treat chronic lymphocytic leukemia (CLL) and multiple sclerosis. Anti-Human CD52 (Alemtuzumab) utilizes the same variable regions from the therapeutic antibody Alemtuzumab making it ideal for research projects.

### Product Info

<b>Amount :</b>	100 µg
<b>Purification :</b>	>95% by SDS-PAGE and HPLC Concentration : $\geq 5.0$ mg/ml
<b>Content :</b>	This biosimilar antibody is aseptically packaged and formulated in 0.01 M phosphate buffered saline (150 mM NaCl) PBS pH 7.2 - 7.4 with no carrier protein, potassium, calcium or preservatives added.
<b>Storage condition :</b>	Functional grade biosimilar antibodies may be stored sterile as received at 2-8°C for up to one month. For longer term storage, aseptically aliquot in working volumes without diluting and store at -80°C. Avoid Repeated Freeze Thaw Cycles.

### Application Note

Endotoxin :  $\leq 1.0$  EU/mg as determined by the LAL method

The suggested concentration for Alemtuzumab biosimilar antibody for staining cells in flow cytometry is  $\leq 0.25$  µg per 10<sup>6</sup> cells in a volume of 100 µl. Titration of the reagent is recommended for optimal performance for each application.