

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 12-8078: Anti-Human IL 12/23 (Briakinumab) - Fc Muted™ HRP

Clone Name: ABT-874
Application: ELISA
Reactivity: Human

Alternative Name: IL-12p40; Interleukin 12; Interleukin 23; IL12; IL23; IL-12; IL-23

**Isotype:** Human IgG1lambda

**Immunogen Information:** This antibody was produced by phage display technology.

## **Description**

Expression Host: HEK-293

This non-therapeutic biosimilar antibody uses the same variable region sequence as the therapeutic antibody Briakinumab. Briakinumab recognizes both human IL12 and IL23 via IL-12/23p40. This product is for research use only.

Briakinumab is a human monoclonal antibody targets the p40 subunit shared by interleukins 12 and 23. IL-12 associates with IL-23alpha to form the heterodimeric cytokine IL-23. IL-23 is associated with various autoimmune inflammatory diseases, and is particularly highly expressed in psoriasis skin lesions. In addition, IL-23 is suspected to play a role in tumorigenesis. Briakinumab binds to and neutralizes human IL-12 and IL-23 (via their shared p40 subunit) and is being investigated for the treatment of rheumatoid arthritis, inflammatory bowel disease, and multiple sclerosis. Anti-Human IL 12/23 (Briakinumab) utilizes the same variable regions from the therapeutic antibody Briakinumab making it ideal for research projects.

## **Product Info**

Amount:  $100 \mu g$ 

Concentration: 0.5 mg/ml

Content: This HRP-conjugated antibody is formulated in 0.01 M phosphate buffered saline (150 mM NaCl)

PBS pH 7.2 - 7.4, 1% BSA. (Warning: Use of sodium azide as a preservative will inhibit the

enzyme activity of horseradish peroxidase)

**Storage condition :** This horseradish peroxidase conjugated monoclonal antibody is stable when stored at 2-8°C. Do

not freeze.

## **Application Note**

The suggested concentration for Briakinumab biosimilar antibody for staining cells in flow cytometry is  $\leq$  1.0  $\mu$ g per 106 cells in a volume of 100  $\mu$ l. Titration of the reagent is recommended for optimal performance for each application. ELISA